arXiv:math/0402155v1 [math.GT] 10 Feb 2004

The Casson-Walker-Lescop invariant of periodic three-manifolds

Nafaa Chbili[1]

Department of Mathematics

Tokyo Institute of Technology

Oh-okayama Meguro Tokyo 152-8551 JAPAN

E-mail: chbili@math.titech.ac.jp

Abstract. Let p be an odd prime and G the finite cyclic group of order p. We use the Casson-Walker-Lescop invariant to find a necessary condition for a three-manifold to have an action of G with a circle as the set of fixed points.

Key words. Casson-Walker-Lescop invariant, group actions on 3-manifolds, global surgery formula, Conway polynomial.

AMS Classification. 57M27, 57M25.

I- INTRODUCTION

In 1985, A. Casson introduced an integer valued invariant of oriented integral homology spheres which reduces modulo two to the Rokhlin invariant. In some sens, the Casson invariant counts the number of conjugacy classes of representations of the fundamental group into $SU(2)$. K. Walker [33] extended the Casson invariant to rational homology spheres and established a method of defining this invariant, that relies only on the surgery presentation of the considered manifold. By a purely combinatorial approach, C. Lescop [19] extended the Casson-Walker

---

[1] Supported by a fellowship from the Japan Society for the Promotion of Science (JSPS), and by a Grant-in-Aid for JSPS fellows 03020.

invariant to any oriented 3-manifold. Namely, given a 3-manifold M and L a surgery presentation of M, Lescop introduced an explicit formula for the Casson-Walker invariant defined in terms of the Alexander polynomials, linking numbers and framing coefficients of the sub-links of L. Lescop's formula will be our main tool in this paper.

Let M be an oriented closed 3-manifold and G a finite group. An intriguing question in low dimensional topology is to know whether the group G acts non trivially on M. Many different approaches were applied to study this question. In the last decade, the quantum invariants were used in order to find necessary conditions for a three-manifold to have such an action, see [1], [4], [5], [6] and [10]. The main motivation of this approach is the fact that the quantum invariants of links were used in a successful way to study symmetries of links and knots in the three-sphere.

A link L in S^3 is said to be p-periodic if there exists an orientation-preserving periodic transformation h of order p such that $fix(h) \cong S^1, h(L) = L$ and $fix(h) \cap L = \emptyset$. By the positive solution of the Smith conjecture [2], we may assume that h is a rotation by a $2\pi/p$ angle around the z-axis. An oriented closed 3-manifold is said to be p-periodic if the finite group $\mathbb{Z}/p\mathbb{Z}$ acts semi-freely on M, with a circle as the set of fixed points.

A p-periodic link is said to be strongly $p-$periodic if each component of the quotient link $\overline{L}$ has linking number zero modulo p with the axis of the rotation. A strongly periodic link L is said to be orbitally separated if the quotient link is algebraically split. In [28], Przytycki and Sokolov proved that a three-manifold is p-periodic if and only if M is obtained from the three-sphere by surgery along a strongly p-periodic link.

There are several studies about the Casson invariant of cyclic branched covers. In particular, Davidow [7], Hoste [15] and Ishibe [16], studied independently the case of Whitehead doubles of knots. Mullins [25], considered the case of 2-fold branched covers along links and provided a nice formula relating the Casson-Walker invariant to the 2-signature of the link and some derivatives of the Jones polynomial. In [9], Garoufalidis and Kricker computed the LMO invariant of cyclic branched covers and gave a formula for the Casson-Walker invariant under the assumption that the branched cover is a rational homology three-sphere.

The purpose of the present paper is to study the Casson-Walker-Lescop invariant of periodic three-manifolds. The only restriction we need here is that the quotient manifold is an integral homology three-sphere. In other words, we deal with manifolds which arise as cyclic branched covers, along knots, of integral homology spheres.

We first study the second coefficient of the Conway polynomial of strongly periodic links. Then we use the surgery formula introduced by Lescop to find a necessary condition for a three-manifold to be periodic.

Notations: Let M be an oriented closed three-manifold. We denote by $\lambda(M)$ the Casson-Walker-Lescop invariant of M. It is worth mentioning that λ takes values in $\frac{1}{12}\mathbb{Z}$. Let $L = l_1 \cup \ldots \cup l_n$ be a $n-$component framed link. For $1 \leq i \leq n$, let l_{ii} be the framing coefficient of the component l_i and define l_{ij} to be the linking number of l_i and l_j if $i \neq j$. Let $E(L) = (l_{ij})$ be the symmetric linking matrix of L. By $Sig(E(L))$ we denote the signature of $E(L)$. Finally, $|H_1(M)|$ denotes the number of elements of the first homology group $H_1(M)$ if finite and zero otherwise. Here are the main results that we will prove in this paper.

Theorem 1. *Let $p \geq 3$ be a prime and M a p-periodic 3-manifold such that the quotient $\overline{M}$ is an integral homology sphere. Then there exists an orbitally separated strongly p-periodic link such that M is obtained from S^3 by surgery along L and :*

$$24\lambda(M) \equiv 3|H_1(M)|Sig(E(L)) \text{ modulo } p,$$

Corollary 1. *Let $p \geq 3$ be a prime and M a $p-$periodic manifold. If the first Betti number of M is not zero, then $24\lambda(M) \equiv 0$ modulo p.*

Let M be an integral homology sphere. We know that M can be obtained from S^3 by surgery along an algebraically split link. On the other hand, we know that if M is p-periodic then M has a strongly p-periodic orbitally separated surgery presentation. One may ask if a periodic homology sphere can be obtained from S^3 by surgery along an algebraically split strongly periodic link.

Corollary 2. *Let $p \geq 3$ be a prime and M a p-periodic 3-manifold. If M has a strongly p-periodic algebraically split surgery presentation then $24\lambda(M) \equiv 0$ modulo p.*

Example. Let $T^3 = S^1 \times S^1 \times S^1$ be the three-dimensional torus. The Casson invariant of T^3 is equal to one. By applying Corollary 1, we can see easily that T^3 is not p-periodic for all prime $p > 3$. Same conclusion for the manifold $S^1 \times S^2$.

If L is a link with n components, let ∇_L denote the Conway polynomial (see paragraph 3) of L, we know that $\nabla_L(z) = z^{n-1}(a_0 + a_1 z^2 + \ldots + a_m z^{2m})$, where coefficients a_i are integers. The proof of Theorem 1 relies heavily on the following theorem, which will be proved in paragraph 5.

Theorem 2. *Let $p \geq 3$ a prime and L an orbitally separated strongly p-periodic link. Then the coefficient a_1 is null modulo p.*

By considering the trefoil knot which is 3-periodic, we can see easily that the condition given by the Theorem 2 is not satisfied by periodic links in general. Indeed, a simple computation shows that for the trefoil knot the coefficient $a_1 = 1$, hence it is not null modulo 3.

This paper is outlined as follows. In section 2, we introduce strongly periodic links and some of their properties. In section 3, we review the surgery presentation of periodic manifolds. Section 4 is to review some properties of the Conway polynomial needed in the sequel. The proof of Theorem 2 is given in section 5. In section 6, we briefly recall Lescop's surgery formula. Ultimately, in section 7, we give the proof of Theorem 1.

II- STRONGLY PERIODIC LINKS.

In this paragraph we recall some definitions.

Definition 2.1. *Let $p \geq 2$ be an integer. A link L of S^3 is said to be p-periodic if and only if there exists an orientation-preserving auto-diffeomorphism of S^3 such that:*

1- Fix(h) is homeomorphic to the circle S^1,

2- the link L is disjoint from Fix(h),

3- h is of order p,

4- $h(L) = L$.

If L is periodic we will denote the quotient link by $\overline{L}$.

The standard example of such a diffeomorphism is given as follows. Let us consider S^3 as the sub-manifold of $\mathbb{C}^2$ defined by $S^3 = \{(z_1, z_2) \in \mathbb{C}^2; |z_1|^2 + |z_2|^2 = 1\}$ and φ_p the following diffeomorphism:

$$\begin{array}{cccc} \varphi_p : & S^3 & \longrightarrow & S^3 \\ & (z_1, z_2) & \longmapsto & (e^{\frac{2i\pi}{p}} z_1, z_2). \end{array}$$

The set of fixed points for φ_p is the circle $\Delta = \{(0, z_2) \in \mathbb{C}^2; |z_2|^2 = 1\}$. If we identify S^3 with $\mathbb{R}^3 \cup \infty$, Δ may be seen as the standard z-axis. By the positive solution of the Smith conjecture [2], a periodic diffeomorphism such as in the previous definition is conjugate to the standard rotation φ_p. Consequently, if L is a p-periodic link then there exists a tangle T such that L is the closure of T^p.

Recall here that if the quotient link $\overline{L}$ is a knot then the link L may have more than one component. In general, the number of components of L depends on the linking numbers of components of $\overline{L}$ with the axis of the rotation.

Recently, Przytycki and Sokolov [28] introduced the notion of strongly periodic links as follows.

Definition 2.2. *Let $p \geq 2$ be an integer. A p-periodic link L is said to be strongly p-periodic if and only if one of the following conditions holds:*

i) The linking number of each component of $\overline{L}$ with the axis Δ is congruent to zero modulo p.

ii) The group $\mathbb{Z}/p\mathbb{Z}$ acts freely on the set of components of L.

iii) The number of components of L is p times greater than the number of components $\overline{L}$.

Remark 2.1. According to condition iii) in the previous definition, a p-strongly periodic link L has $p\alpha$ components, where α is the number of components of the quotient link $\overline{L}$. These $p\alpha$ components are divided into α orbits with respect to the free action of $\mathbb{Z}/p\mathbb{Z}$ (condition i)). Assume that $\overline{L} = l_1 \cup \ldots \cup l_\alpha$, there is a natural cyclic order on the components of L. Namely,

$$L = l_1^1 \cup \ldots \cup l_1^p \cup l_2^1 \cup \ldots \cup l_2^p \cup \ldots \cup l_\alpha^1 \cup \ldots \cup l_\alpha^p$$

where $\varphi_p(l_i^t) = l_i^{t+1}\ \forall 1 \leq t \leq p-1$ and $\varphi_p(l_i^p) = l_i^1$, for all $1 \leq i \leq \alpha$.

Example. The link in the following figure is a strongly 5-periodic link.



Figure 1.

Definition 2.3. *Let L be a link in the three-sphere. L is said to be algebraically split if and only if the linking number of any two components of L is null.*

Definition 2.4. *Let $p \geq 2$ be an integer. A strongly p-periodic link is said to be orbitally separated if and only if the quotient link is algebraically split.*

Remark 2.2. Throughout the rest of this paper we use the term OS link to refer to an orbitally separated link. If L is a strongly p-periodic OS link, then for all s and $i \neq j$ we have $\sum_{t=1}^{p} lk(l_i^s, l_j^t) = 0$.

III- PERIODIC 3-MANIFOLDS

Let M be a 3-manifold. It is well known from Lickorish's work [21] that M may be obtained from the three-sphere S^3 by surgery along a framed link L in S^3. Such a link is called a surgery presentation of M. This section deals with surgery presentations of periodic three-manifolds.

Definition 3.1. *Let $p \geq 2$ be an integer. A 3-manifold M is said to be p-periodic if and only if there exists an orientation-preserving auto-diffeomorphism of M, such that:*

1- Fix(h) is homeomorphic to the circle S^1,

2- h is of order p.

The three-sphere S^3 is p-periodic for all $p \geq 2$. The acting diffeomorphism is nothing but the rotation φ_p defined in section 2. Another concrete example is the Brieskorn manifold $\Sigma(p,q,r)$, which has periods p, q and r. One can easily give explicit formulas for the corresponding diffeomorphisms by considering $\Sigma(p,q,r)$ as the intersection of the complex surface $\{(z_1, z_2, z_3) \in \mathbb{C}^3; z_1^p + z_2^q + z_3^r = 0\}$ with an appropriate sphere of dimension 5 (see [23]).

Theorem 3.2 ([28]). *Let p be a prime. A 3-manifold M is p-periodic if and only if M is obtained from S^3 by surgery along a strongly $p-$periodic link.*

Remark. If the quotient manifold is a homology sphere then we may suppose that L is orbitally separated.

IV- THE CONWAY POLYNOMIAL

The Alexander polynomial of periodic knots has been subject of an extensive literature. In 1971, Murasugi [25] proved a congruence relationship between the Alexander invariant of a periodic knot K and the Alexander invariant of the quotient knot. Later, different approaches were used to give alternative proofs to Murasugi's result [3], [8], [12]. In [30] and [22], similar results were proved for the multi-variable Alexander polynomial. Murasugi's result was applied to rule out the possibility of being periodic for many knots.

In this paper we consider the Conway version of the Alexander polynomial. The Conway polynomial ∇ is an invariant of ambient isotopy of oriented links which can be defined uniquely by the following:

$$\textbf{(i)} \quad \nabla_{\bigcirc}(z) = 1$$

$$\textbf{(ii)} \quad \nabla_{L_+}(z) - \nabla_{L_-}(z) = z\nabla_{L_0}(z),$$

where $\bigcirc$ is the trivial knot, L_+, L_- and L_0 are three oriented links which are identical except near one crossing where they look like in the following figure:



FIGURE 2

It is well known that if L is a link with n components then the Conway polynomial of L is of the form $\nabla_L(z) = z^{n-1}(a_0 + a_1z^2 + \ldots + a_mz^{2m})$, where coefficients a_i are integers. The first coefficient depends only on the linking matrix of the link. The second coefficient a_1 is the first non trivial Vassiliev invariant for knots. It is related to the Casson-Walker invariant by the global surgery formula introduced by Lescop [19].

At the beginning, let us recall some properties of the Conway polynomial needed in the sequel. Let $L = l_1 \cup \ldots \cup l_n$ be an oriented n component link in the three-sphere. Let $l_{ij} = \mathrm{Lk}(l_i, l_j)$ if $i \neq j$ and define $l_{ii} = -\sum_{j=1, j\neq i}^{n} l_{ij}$. Define the linking matrix $\mathcal{L}$ as $\mathcal{L} = (l_{ij})$. The matrix $\mathcal{L}$ is symmetric and each row adds to zero. Hence, every cofactor of this matrix is the same. We refer the reader to [11], [13] and [14], for details about the linking matrix.

Theorem 4.3 [14]. *Let L an oriented link with n components. Then $a_0 = \mathcal{L}_{ij}$, where $\mathcal{L}_{ij}$ is any cofactor of the the linking matrix $\mathcal{L}$.*

It follows from this theorem, that if L is algebraically split then the first coefficient of the Conway polynomial is zero. Levine [20] proved that the coefficient of z^i is null, for all $i \leq 2n-3$, and gave an explicit formula for the coefficient of z^{2n-2}.

Proposition 4.4 [20]. *If L is algebraically split with n components, then $\nabla_L(z)$ is divisible by z^{2n-2}.*

Let Σ_1, Σ_2 and Σ_3 be three oriented surfaces imbedded in S^3. For $1 \leq i \leq 3$, let n_i be a positive normal vector to Σ_i. Assume that the three surfaces intersect transversally in their interiors. Let $A = \{a_1, \ldots, a_s\} = \Sigma_1 \cap \Sigma_2 \cap \Sigma_3$, every $a_i \in A$ may be occupied with a sign $\epsilon(a_i) = \pm 1$, depending on the orientation of the basis defined by the three normal vectors. Let $\Sigma_1 : \Sigma_2 : \Sigma_3 = \sum_{i=1}^{s} \epsilon(a_i)$.

Let $L = l_1 \cup \ldots \cup l_n$ be an algebraically split link. We can choose Seifert surfaces Σ_1, ... , Σ_n such that l_i bounds Σ_i and l_j does not intersect Σ_i if $i \neq j$. We define $\bar{\mu}_{ijk}$ to be $\Sigma_1 : \Sigma_2 : \Sigma_3$ if i, j and k are distinct and to be zero otherwise. Now we define u to be the $(n-1)\times(n-1)$-matrix defined by $u_{ij} = \sum_{k=1}^{n} \bar{\mu}_{ijk}$, for every $1 \leq i, j \leq n-1$. Levine [20], proved the following.

Theorem 4.5 [20]. *If $L = l_1 \cup \ldots \cup l_n$ is an algebraically split link then: $a_{n-1} = det(u)$.*

V- PROOF OF THEOREM 2

This paragraph is devoted to the proof of Theorem 2. Let L be a strongly p-periodic link in the three-sphere. Let $\overline{L}$ be the factor link, so here we have $L = \pi^{-1}(\overline{L})$, where π is the canonical surjection corresponding to the action of the rotation on the three-sphere. Let $\overline{L}_+$, $\overline{L}_-$ and $\overline{L}_0$ denote the three links which are identical to $\overline{L}$ except near one crossing where they are like in figure 2. Now, let $L_{p+} := \pi^{-1}(\overline{L}_+)$, $L_{p-} := \pi^{-1}(\overline{L}_-)$ and $L_{p0} := \pi^{-1}(\overline{L}_0)$.

Lemma 5.1. *Let p be a prime, then we have the following congruence modulo p:*

$$\nabla_{L_{p+}}(z) - \nabla_{L_{p-}}(z) \equiv z^p \nabla_{L_{p0}}(z).$$

Proof. Przytycki [26] proved a similar lemma for the skein (HOMFLY) polynomial of periodic links. The same argument can be used here to prove our lemma.

Notice that if L_{p+} is a strongly p-periodic link then L_{p-} is also a strongly periodic link. However, two cases are to be considered for the link L_{p0}.

First case: if the considered crossing of the factor link is a mixed crossing, (involving two different components of the factor link say K_1 and K_2). After the crossing change, the two components connect and give birth to a new component K_{12}, as the linking numbers of K_1 and K_2 with the rotation axis is null modulo p then the linking number of K_{12} with the rotation axis is also null modulo p as it is the sum of two zeros. Consequently, the link L_{p0} is strongly p-periodic.

Second case: If the considered crossing is a self crossing, in this case the link L_{p0} may be not strongly p-periodic.

Let L be a strongly p-periodic link, then we know that L has $p\alpha$ components, where α is a positive integer. The proof of Theorem 2, will be done by induction on α. Recall here that α is the number of components of the factor link $\overline{L}$.

It is easy to see that each of L_{p+} and L_{p-} has $p\alpha$ components. Let s be the number of components of the link L_0. Let

$\nabla_{L_{p+}}(z) = z^{p\alpha-1}(a_0 + a_1 z^2 + \ldots + a_m z^{2m})$,

$\nabla_{L_{p-}}(z) = z^{p\alpha-1}(b_0 + b_1 z^2 + \ldots + b_k z^{2k})$,

$\nabla_{L_{p0}}(z) = z^{s-1}(c_0 + c_1 z^2 + \ldots + c_r z^{2r})$.

Lemma 5.2. *Let $p \geq 3$ be a prime and L a strongly p-periodic link. Assume that L has p components, then:* $a_1(L) \equiv 0$ modulo p.

Proof. Here $\alpha = 1$, then each of L_{p+} and L_{p-} has p components. By lemma 5.1, we have the following congruence modulo p:

$$z^{p-1}(a_0 + a_1 z^2 + \ldots + a_m z^{2m}) - z^{p-1}(b_0 + b_1 z^2 + \ldots + b_k z^{2k}) \equiv z^{p+s-1}(c_0 + c_1 z^2 + \ldots + c_r z^{2r}).$$

From this identity we conclude that if $s \neq 2$ then $a_1 - b_1 \equiv 0$ modulo p. If $s = 2$, then $a_1 - b_1 \equiv c_0$ modulo p.

If L_{p0} has two components then by the Hoste Formula we know that c_0 is minus the linking number of the two components. As the link $\overline{L}_0$ has two components (because $\overline{L}_+$ is a knot), then the linking number of the two components of L_{p0} is p times the linking number of the two components of the link $\overline{L}_{p0}$. Hence c_0 is zero modulo p.

Consequently, L_{p+} and L_{p-} have the same a_1 coefficient modulo p. Notice, that starting from a strongly $p-$periodic link L, one may change crossings along orbits to get an algebraically split link. The algorithm can be described as follows. We start by changing the crossings of l_1^1 and l_1^2, until having their linking number zero. Of course, at the same moment we get the same for l_1^2 and l_1^3 ... etc. In the second step we do the same for l_1^1 and l_1^3. Obviously by this procedure we get an algebraically split link. If $p > 3$, then using proposition 4.4 we conclude that $a_1(L)$ is zero modulo p. It remains to check the case $p = 3$.

For $1 \leq i \leq 3$, let $\sum_i$ be a Seifert surface which is bounded by the component l_1^i. Following the notations in paragraph 4, recall that $\bar{\mu}_{ijk}$ is defined to be the algebraic sum of the triple points of $\sum_i \cap \sum_j \cap \sum_k$ if i, j and k are distinct and to be zero otherwise. Now let $u_{ij} = \sum_{k=1}^{3} \bar{\mu}_{ijk}$. It is well known [20], that a_1 is the determinant of the matrix u_{ij}. In our case we can choose three Seifert surfaces such that $\varphi(\sum_1) = \sum_2$, $\varphi(\sum_2) = \sum_3$ and $\varphi(\sum_3) = \sum_1$. Assume that $M \in \sum_i \cap \sum_j \cap \sum_k$, then $\varphi(M)$ and $\varphi^2(M)$ are also in the intersection of the three surfaces. If $M \notin \Delta$ then the three points are distinct and the sum gives zero modulo three. If $M \in \Delta$, then as the linking number of the factor knot with the axis is zero modulo three, then the number of intersections points (with signs) of a the quotient surface $\overline{\sum}$ and the axis Δ is a multiple of 3. It is not hard to see that the matrix u_{ij}, is null modulo 3. Hence the coefficient a_1 is null modulo 3.

Suppose now that Theorem 2 is true for strongly periodic OS links with a number of components less than or equal to $p\alpha$. Let L be a strongly p-periodic OS link having $p(\alpha+1)$ components. It is easy to see that we have essentially two cases.
Case 1: If we change a mixed crossing of the factor link then the resulting link L_{p0} will be a strongly p-periodic OS link with $p\alpha$ components. If we write lemma 5.1 in this case we can see that:

$$z^{p(\alpha+1)-1}(a_0+a_1z^2+\ldots+a_mz^{2m})-z^{p(\alpha+1)-1}(b_0+b_1z^2+\ldots+b_kz^{2k}) \equiv z^{p+p\alpha-1}(c_0+c_1z^2+\ldots+c_rz^{2r}).$$

Hence, $a_1 - b_1 \equiv c_1$ modulo p. By the induction hypothesis the coefficient c_1 is zero modulo p.

Case 2: If we change a self crossing of the factor link. There are two possible situations:
First, If the resulting link L_{p0} has more components than L_{p+}, in this case we can see immediately from lemma 5.1 that $a_1 \equiv b_1$ modulo p.

Second, if the resulting link L_{p0} has $p\alpha+2$ components. In this case only one orbit is affected and the link L_{p0} is made up of two invariant components K_1 and K_2 and a strongly p-periodic link with $p\alpha$ components.

$$L_{p0} = K_1 \cup K_2 \cup l_2^1 \cup \ldots \cup l_2^p \cup l_3^1 \cup \ldots \cup l_3^p \ldots \cup l_{(\alpha+1)}^1 \cup \ldots \cup l_{(\alpha+1)}^p.$$

From lemma 5.1 we have:

$$z^{p(\alpha+1)-1}(a_0+a_1z^2+\ldots+a_mz^{2m})-z^{p(\alpha+1)-1}(b_0+b_1z^2+\ldots+b_kz^{2k}) \equiv z^{p+p\alpha+1}(c_0+c_1z^2+\ldots+c_rz^{2r}).$$

Consequently, $a_1 - b_1 \equiv c_0$ modulo p. It remains now to prove that the coefficient c_0 vanishes modulo p.

The coefficient c_0 can be computed using the Hoste formula [14]. Let us now compute the coefficient c_0 in the case of our link L_0. Recall that we assumed that we only changed the first orbit $l_1^1 \cup \ldots \cup l_1^p$ of the link L. This sub-link is transformed onto the link $K_1 \cup K_2$. It can be easily seen that each of K_1 and K_2 is invariant by the rotation. Let $\overline{K}_1$ and $\overline{K}_2$ the corresponding factor knots. We have the following $Lk(K_1, K_2) = pLk(\overline{K}_1, \overline{K}_2) \equiv 0$ modulo p and

$$Lk(K_i, l_j^1) = Lk(K_i, l_j^2) = \ldots = Lk(K_i, l_j^p) \ \ \forall \ \ 1 \leq i \leq 2 \text{ and } 2 \leq j \leq \alpha + 1.$$

Furthermore, as the link L is orbitally separated we have $\sum\limits_{t=1}^{p} Lk(l_1^i, l_j^t) = 0$ for all i and all $j \neq 1$. Thus, $\sum\limits_{t=1}^{p} Lk(K_1, l_j^t) + \sum\limits_{t=1}^{p} Lk(K_1, l_j^t) = 0$. From this we conclude that for all $2 \leq t \leq \alpha + 1$ we have $pLk(K_1, l_j^t) + pLk(K_2, l_j^t)$ is zero. Hence, $Lk(K_1, l_j^t) = -Lk(K_2, l_j^t)$. Consequently the linking matrix of L_{p0} is of the form (here coefficients are considered modulo p).

$$\begin{pmatrix}
0 & 0 & t_2^1 & t_2^2 & \ldots & t_{\alpha+1}^p \\
0 & 0 & -t_2^1 & -t_2^2 & \ldots & -t_{\alpha+1}^p \\
t_2^1 & -t_2^2 & . & . & \ldots & . \\
t_2^2 & -t_2^2 & . & . & \ldots & . \\
. & . & . & . & \ldots & . \\
. & . & . & . & \ldots & . \\
. & . & . & . & \ldots & . \\
t_{\alpha+1}^p & -t_{\alpha+1}^p & . & . & \ldots & .
\end{pmatrix}$$

We know that in the linking matrix of a link the sum of lines is zero. Moreover in our matrix the first line and the second line are dependant. Thus, all cofactors of the matrix are zero. Consequently, L_{p+} and L_{p-} have the same a_1 coefficient modulo p.

Remark. *As explained above, starting from a strongly p-periodic OS link L_{p+}, one may change crossings along an orbit without modifying the coefficient a_1 modulo p. It is worth mentioning that to prove this result we only used the fact that the link L_{p0} is OS. Hence, we* *may apply the same procedure again to the link L_{p-}, although L_{p-} is not orbitally separated.*

In order to transform the link L onto an algebraically split link, we apply the same techniques as in the case $\alpha = 1$. Obviously by this procedure all orbits will be made up of algebraically split sub-links. Now, we apply the same techniques for components belonging to different orbits until getting an algebraically split link. Finally, by using Proposition 4.4 we conclude that $a_1(L)$ is zero modulo p. This ends the proof of Theorem 2.

VI- LESCOP'S SURGERY FORMULA

In [19], Lescop provided a combinatorial description of the Casson-Walker invariant. Indeed, Lescop introduced three equivalent formulas relating the Casson-Walker invariant to the Conway polynomial and the multi-variable Alexander polynomial. This allowed her to extend the Casson-Walker invariant to all oriented three-manifolds. In this paragraph we roughly recall some notations, we refer the reader to [19], (page 19 and 20) for more details.

Throughout the rest of this paper, $N = \{1, \dots, n\}$. Let $L = \cup_1^n l_i$ an n component link in the three-sphere. For every $I \subset N$, we define $L_I = \cup_{i \in I} l_i$. Let G be a graph with n vertices $\{v_1, \dots, v_n\}$ and e_{ij} the edge of G with end points v_i and v_j. The linking number of L with respect to the edge e_{ij}, $\mathrm{Lk}(L, e_{ij})$ is defined to be the linking number of l_i and l_j. The linking of L with respect to the graph G is defined by:

$$\mathrm{Lk}(L, G) = \prod_{e \text{ edge of G}} \mathrm{Lk}(L, e).$$

Let $\mathcal{C}$ be the set of graphs G with n vertices such that G is topologically an oriented circle. We define the circular linking number of the link L by the following:

$$\mathrm{Lk}_c(L) = \sum_{C \in \mathcal{C}} \mathrm{Lk}(L, C).$$

Now we define the integer θ_b as follows:

$$\theta_b(L_I) = \sum_{K\subset I} \mathrm{Lk}_c(L_K) \sum_{(i,j)\in K^2} \sum_{g\in\sigma_{I\setminus K}} l_{ig(1)} l_{g(1)g(2)} \ldots . l_{g(\sharp(I\setminus K))j}$$

where $\sigma_{I\setminus K}$ denotes the set of bijections from $\{1,\ldots,\sharp I\setminus K\}$ to $I\setminus K$. Recall here that we consider surgery with integral coefficients, hence the formula 1.7.6 in [19] can be written as follows

$$\theta(L_I) = \begin{cases} \theta_b(L_I) & \text{if } I > 2 \\ \theta_b(L_I) - 2l_{ij} & \text{if } I = \{i,j\} \\ \theta_b(L_I) + 2 & \text{if } I = \{i\} \end{cases}$$

If $I \subset \{1,2,\ldots,n\}$, we define $E(L_I)$ as the linking matrix of L_I and we define $E(L_{N\setminus I}; I) = (l_{ijI})_{i,j\in N\setminus I}$ where:

$$l_{ijI} = \begin{cases} l_{ij} & \text{if } i \neq j \\ l_{ii} + \sum_{k\in I} l_{ik} & \text{if } i = j \end{cases}$$

Let M be a three manifold obtained from S^3 by integral surgery along a framed link L in S^3. Let $E(L)$ be the symmetric linking matrix of L. Let $b_-(L)$ be the number of negative eigenvalues of $E(L)$ and define $sign(L)$ to be $(-1)^{b_-(L)}$. With respect to this notations, the Casson-Walker invariant of M is given by the following formula.

$$\begin{aligned}\lambda(M) = \quad & sign(L) \sum_{J\subset N/J\neq\emptyset} det(E(L_{N\setminus J}, J)) a_1(L_J) \qquad + \\ & sign(L) \sum_{J\subset N/J\neq\emptyset} \frac{det(E(L_{N\setminus J}))(-1)^{\sharp J}\theta(L_J)}{24} \quad + \\ & |H_1(M)| \frac{Sig(E(L))}{8}.\end{aligned}$$

This formula corresponds to Proposition 1.7.8 in [19]. As we only consider integral surgery then for all i we have $q_i = 1$ and the Dedekind sums are null.

VII- PROOF OF THE MAIN THEOREM

Let M be a p-periodic 3-manifold and L a strongly p-periodic surgery presentation of L. Let:

$$D_1 = sign(L) \sum_{J \subset N / J \neq \emptyset} det(E(L_{N \setminus J}, J)) a_1(L_J)$$
$$D_2 = sign(L) \sum_{J \subset N / J \neq \emptyset} \frac{det(E(L_{N \setminus J}))(-1)^{\sharp J} \theta(L_J)}{24}.$$

Lemma 7.1. *Let p be an odd prime and L an OS strongly p-periodic link. Then D_1 is null modulo p.*

Proof. The summation is done over all sub-links of L. It is easy to see that if the sub-link L_J is not periodic then there is p sub-links of L which are the same as L_J. Hence, the contributions of these links add to zero modulo p. However if the sub-link is p-periodic, it is necessary an OS strongly p-periodic link. By the Theorem 2, the second coefficient of the Conway polynomial of L_J is null modulo p. This ends the proof of lemma 7.1.

Lemma 7.2. *Let p be an odd prime and L an OS strongly p-periodic link. Then $24D_2$ is null modulo p.*

Proof. In the term D_2, the sum is taken over all sub-links of L. As in the proof of the previous lemma, the contributions of non periodic sub-links add to zero modulo p. Hence, we have to consider only strongly p-periodic sub-links. Recall that if L_I is a link with at least 3 components, then

$$\theta(L_I) = \theta_b(L_I) = \sum_{K \subset I} \mathrm{Lk}_c(L_K) \sum_{(i,j) \in K^2} \sum_{g \in \sigma_{I \setminus K}} l_{ig(1)} l_{g(1)g(2)} \ldots . l_{g(\sharp(I \setminus K))j}.$$

Let L_J a periodic sub-link of L. Here also, for the computation of $\theta(L_J)$ modulo p, we need to consider only periodic sub-links of L_J. If L_K is a periodic sub-link, then the term $l_{ig(1)} l_{g(1)g(2)} \ldots . l_{g(\sharp(I \setminus K))j}$ does not change if the indexes are cyclically permuted. Thus, we conclude that $24D_2$ is null modulo p. This ends the proof of Lemma 7.2.

The proof of Corollary 2 is based on the fact that if L is an algebraically split p-strongly periodic then the linking matrix of L is diagonal and each diagonal coefficient of the matrix appears exactly sp times, for some integer s. Hence, the signature of $E(L)$ is null modulo p. The proof of Corollary 1 is obvious.

References

[1] J. K. Bartoszynska, P. Gilmer and J. Przytycki. *3-Manifold invariants and periodicity of homology spheres.* Algebraic and Geometric Topology, Volume 2 pp. 825-842, (2002).

[2] H. Bass and J. W. Morgan. *The Smith conjecture.* Pure and App. Math. 112, New York Academic Press (1994).

[3] G. Burde. *Über periodische knoten.* Arkiv der math. (Basel) 30, pp. 487-492, 1978.

[4] N. Chbili. *Les invariants θ_p des 3-variétés périodiques.* Annales de l'institut Fourier, Fascicule 4, pp. 1135-1150 (2001).

[5] N. Chbili. *Quantum invariants and finite group actions on 3-manifolds*, Topology Appl. vol 136/1-3 pp. 219-231, 2004.

[6] Q. Chen and T. Le. *Quantum invariants of periodic links and periodic 3-manifolds.* Preprint.

[7] A. Davidow. *Casson's invariant and twisted double knots.* Topology Appl. 58 pp. 93-101, 1994.

[8] J. F. Davis, C. Livingston. *Alexander polynomials of periodic knots.* Topology, 30, pp. 551-564, 1991.

[9] S. Garoufalidis and A. Kricker. *Finite type invariants of Cyclic branched covers*, Preprint.

[10] P. Gilmer. *Quantum invariants of periodic three-manifolds.* Geometry and Topology monographs, Vol. 2: proceeding of the Kirbyfest, pp. 157-175 (1999).

[11] R. Hartley. *The Conway potential function for links.* Comment. Math. Helv. 58 (1983), no. 3, 365–378.

[12] J. Hillman. *New proofs of two theorems on periodic knots.* Archiv. Math. 37, pp. 457-461, 1981.

[13] F. Hosokawa. *On ∇-polynomials of links.* Osaka Math. J. 10 1958 273–282. d

[14] J.Hoste. *The first coefficient of the Conway polynomial.* Proc. Amer. Math. Soc., 95, pp. 299-302, 1985

[15] J.Hoste. *A formula for Casson's invariant.* Trans. Amer. Math. Soc.,297, pp. 547-562, 1986

[16] K. Ishibe. *The Casson-Walker invariantfor branched cyclic coversof S^3 branched over a doubled knot.* Osaka J. Math. 34, pp. 481-495, 1997.

[17] V. F. R. Jones. *Hecke algebra representations of braid groups and link polynomials.* Annals of Mathematics, Vol. 126, pp. 335-388, 1987.

[18] R. Kirby. *The calculus of framed links in S^3*, Invent. Math. 45 pp. 35-56 (1978).

[19] C. Lescop. *Global surgery formula for the Casson-Walker invariant*, Annals of Mathematics Studies, Princeton Univ. Press (1996).

[20] J. Levine. *The Conway polynomial of an algebraically split link.* Proceeding of knots 96, edited by S. Suzuki, World Scientific Publishing Co, pp. 23-29, 1997.

[21] W. B. R. Lickorish. *A representation of orientable combinatorial 3-manifolds.* Ann. Math. 76 (1962), 531-540.

[22] Y. Miyazawa. *Conway polynomials of periodic links.* Osaka J. math. 31, pp. 147-163, 1994.

[23] J. Milnor. *Singular points and complex hypersurfaces.* Ann. Math. studies, Princeton University Press. 1968.

[24] D. Mullins. *The generalized Casson invariant for 2-fold branched covers of S^3 and the Jones polynomial.* Topology, 32 419-438, 1993.

[25] K. Murasugi. *On periodic knots.* Comment. Math. Helv. 46, pp. 162-174, 1971.

[26] J. H. Przytycki. *On Murasugi's and Traczyk's criteria for periodic links.* Math. Ann., 283, pp. 465-478, 1989

[27] J. H. Przytycki. *An elementary proof of the Traczyk-Yokota criteria for periodic knots.* Proc. Amer. Math. Soc., 123, pp. 1607-1611, 1995.

[28] J. Przytycki and M. Sokolov. *Surgeries on periodic links and homology of periodic 3-manifolds.* Math. Proc. Cambridge Phil. Soc. Vol. 131(2), pp 295–307. 2001.

[29] D. Rolfsen. *Knots and Links.* Mathematics Lecture Series, Publish or Perish, Inc. Houston, Texas, 1990.

[30] M. Sakuma. *On the polynomials of periodic links.* Math. Ann. 257, pp. 487-494, 1981.

[31] P. Traczyk. *A criterion for knots of period 3.* Topology appl, 36, pp. 275-281, 1990.

[32] P. Traczyk. *Periodic knots and the skein polynomial.* Invent. Math, 106(1), pp. 73-84, 1991.

[33] K. Walker. *An extension of Casson's invariant.* Ann. of Math. Studies 126, Princeton University Press, 1992.

[34] Y.Yokota. *The skein polynomial of periodic knots.* Math. Ann. 291, pp. 281-291, 1991.